



Contents

- Carone G., Marra A.C. & Mesiano C. – First record of Dugongidae (Mammalia: Sirenia) from the Floresta Calcareous Formation (Late Burdigalian – Early Langhian, Reggio Calabria, Southern Italy)
- Selden P. A., Dunlop J. A. & Simonetto L. – A fossil whip-scorpion (Arachnida: Thelyphonida) from the Upper Carboniferous of the Carnic Alps (Friuli, NE Italy)
- Abbassi N., Domning D. P., Izad Navid N. & Shakeri S. – Sirenia fossils from Qom Formation (Burdigalian) of the Kabudar Ahang area, Northwest Iran.
- Moncunill-Solé B., Tuveri C., Arcrca M. & Angelone C. – Comparing the body mass variations in endemic insular species of the genus *Prolagus* (Ochotonidae, Lagomorpha) in the Pleistocene of Sardinia (Italy).
- Brandano M., Bosellini F. R., Mazzucc hi A. & Tomassetti L. – Coral assemblages and biocostructions adapted to the depositional dynamics of a mixed carbonate siliciclastic setting: the case study of the Burdigalian Bonifacio Basin (South Corsica).
- Monaco P., Famiani F. & La Iacona F. – Bulldozing and resting traces of freshwater mussel *Anodonta woodiana* and substrate characteristics in lake-margin and river settings of Umbria, Italy.
- Bianucci G., Collareta A., Post K., Varola A. & Lambert O. – A new record of *Messapicetus* from the Pietra Leccese (Late Miocene, Southern Italy): antitropical distribution in a fossil beaked whale (Cetacea, Ziphiidae).
- Jan Irfan U., Shah A., Stephenson M. H., Iqbal S., Hanif M., Wagreich M. & Hussain Shahid H. – The sedimentology of the Lower Permian Dandot Formation: a component of the Gondwana deglaciation sequence of the Salt Range, Pakistan.